

Gauge Theory, Geometry and the Large N Limit

Vijay Balasubramanian,^{(1)*}Rajesh Gopakumar^{(1,2)†}and Finn Larsen^{(3)‡}

(1) Lyman Laboratory of Physics, Harvard University, Cambridge, MA 02138

(2) Physics Department, Univ. of California at Santa Barbara, Santa Barbara, CA 93106

(3) David Rittenhouse Laboratories, University of Pennsylvania, Philadelphia, PA 19104

Abstract

We study the relationship between M theory on a nearly lightlike circle and $U(N)$ gauge theory in $p + 1$ dimensions. We define large N limits of these theories in which low energy supergravity is valid. The regularity of these limits implies an infinite series of nonrenormalization theorems for the gauge theory effective action, and the leading large N terms sum to a Born-Infeld form. Compatibility of two different large N limits that describe the same decompactified M theory leads to a conjecture for a relation between two limits of string theories.

1 Introduction

A common theme of recent work in string theory is that conventional geometry should sometimes emerge as an effective low-energy description from an auxiliary gauge theory. This surprising idea emerges in its simplest form from an examination of the low-energy dynamics of parallel D-branes [1, 2]. In this context, the classical geometry of spacetime appears via dynamics on the quantum moduli space of gauge configurations. The connection between gauge theory and geometry has been exploited in M(atric) theory, the conjectured nonperturbative definition of M-theory [3, 4], as well as in the related analysis of M(atric) strings [5]. In these developments, the implied logic is that gauge theory is simpler than string theory, and so constitutes a preferred starting point for the discussion. In the present paper, we pursue the reverse strategy: we consider limits of M-theory where spacetime geometry can be reliably studied in supergravity, and use these to infer properties of gauge theory effective actions.

*vijayb@pauli.harvard.edu

†gopakumr@pauli.harvard.edu

‡larsen@cvtic.hep.upenn.edu

Our starting point is the recent argument by Seiberg that relates M-theory on a nearly lightlike circle to the $U(N)$ gauge theory dynamics of N D-branes [6] (also see [7]). We consider several large N limits where M-theory is adequately described by its low-energy supergravity approximation. We argue that these limits are nonsingular and use the requirement that the corresponding gauge theory limits also be regular to find an infinite series of non-renormalization theorems for the loop expansion of the gauge theory effective action.

Several subtleties arise. For example, a null circle has zero proper length; so low-energy M-theory on a nearly lightlike circle is in danger of breaking down due to short distance effects. However, we find that a large momentum along the circle exerts a pressure that makes it spacelike and large. In our large N limits the M-theory is therefore decompactified in the bulk of spacetime and neither the curvature invariants, nor the proper distance around the circle, approach the 11 dimensional Planck scale. It would be a surprise if 11 dimensional supergravity could not be applied in such regimes where it shows no sign of breaking down.

An important result for the gauge theory is that the leading large N terms to all loops in the gauge theory effective description of graviton scattering sum up to a Born-Infeld form. This result is already known as a prediction of M(atrix)-theory [8, 9, 10, 11, 12], but we recover it using different assumptions. Our large N limits are not the 't Hooft scaling limits which usually make sense for a generic gauge theory. We will comment on how our alternative limits seem to make sense as a consequence of the non-renormalization theorems.

Additionally, we identify certain nonplanar diagrams in the gauge theory with corrections to the 11 dimensional low-energy effective action that are suppressed by powers of the Planck length. We also present a conjecture arising from a comparison between two different large N limits which should, at least naively, yield the same decompactified M theory. This conjecture essentially says that string theory in the large N gauge theory limit is equivalent to the same string theory, but in the rather different domain specified in Sec. 3.3. This equivalence is in a similar spirit (but not exactly the same) as the expected simultaneous validity of supergravity and large N gauge theory in the near horizon limit of black holes [13].

Sec. 2 summarizes the argument that relates M-theory on a nearly lightlike circle and a transverse T^p , to $U(N)$ gauge theory in $p + 1$ dimensions [7, 6] and discusses graviton scattering in terms of effective actions in gauge theory and supergravity. In Sec. 3 we specify the large N limits that allow descriptions in both gravity and gauge theory. The discussion for gravitons is generalized to extended objects and we give some further arguments for the Born-Infeld resummation of the leading contributions to graviton scattering.

2 M-theory and Gauge Theory

2.1 Preliminaries

Consider M-theory on a nearly lightlike circle defined by the identifications $(x_{11}, t) \sim (x_{11} + \sqrt{R^2/2 + R_S^2}, t - R/\sqrt{2})$, where $R_S \ll R$. Also introduce a transverse T^p specified by $r^i \equiv r^i + R^i$.¹ Working in this nearly lightlike frame we consider a probe graviton with longitudinal momentum $P_p^+ = K/R$ impinging on a target graviton with $P_t^+ = N/R$. We will always be interested in situations with $K \ll N$. The probe has small transverse velocity v in a noncompact direction, in the center of mass frame where the heavy target is nearly stationary. The probe and target are separated by b in a noncompact direction other than the direction of motion.

Seiberg's kinematic analysis [6] exploits 11 dimensional Lorentz invariance to apply a boost of rapidity α to the nearly lightlike circle where $\sinh \alpha = R/\sqrt{2}R_S$. Winding modes are important after the boost and can be accounted for by T-dualizing the torus to get a theory of K Dp-brane probes interacting at low energies with a target of N Dp-branes. After further uniform rescaling of all lengths as $\tilde{l} = l(R_S/R)$ the M-theory scattering problem reduces to the dynamics of Dp-branes with effective string coupling and string scale:

$$\tilde{g}_S = \frac{(R_S M_P)^{3/2}}{V_T (R_S M_P^3)^{p/2}} \quad ; \quad \tilde{M}_S^2 = \frac{R^2 M_P^3}{R_S} \quad (1)$$

where $V_T = \prod_i R_i$ is the volume of the original transverse torus in M-theory. The probe branes have a velocity $\tilde{v} = vR_S/R$, kinetic energy $E = Kv^2/R$, and are separated from the target branes by $\tilde{b} = bR_S/R$ where v and b are the original 11 dimensional velocity and separation. The T-dual transverse torus has cycles of sizes $\Sigma_i = 1/R_i R M_P^3$.

With these parameters, for $p \leq 3$, the $R_S \rightarrow 0$ limit yields the standard gauge theory description of the low-energy, short-distance dynamics of Dp-branes. The action for the K probes and N target branes is the dimensional reduction of minimally supersymmetric $U(N+K)$ Yang-Mills in 10 dimensions to $(p+1)$ dimensions:

$$S_0 = \frac{1}{g_{YM}^2} \int d^{p+1}x \frac{1}{4} \text{Tr}(F^2) \quad (2)$$

To perform the dimensional reduction, for $i > p$ the gauge fields A^i are interpreted as scalar fields X^i and all derivatives ∂_i vanish. We will relate the fields A and X to the parameters of the scattering branes in the next section. The coupling g_{YM}^2 (finite as $R_S \rightarrow 0$) is given by:

$$g_{YM}^2 = \frac{\tilde{g}_S}{\tilde{M}_S^{p-3}} = \frac{(R M_P^2)^3}{V_T (R M_P^3)^p} \quad (3)$$

For $p > 3$, the $R_S \rightarrow 0$ limit is more complicated because it leads to large string coupling.

¹We use R^i for the *circumference* of a circle, following many recent works.

2.2 Scattering in gauge theory

The separation of probe from target in M-theory is b and, after the above rescaling of units, the separation between Dp -branes is $\tilde{b} = bR_S/R$ in the boosted theory. The positions of branes in the gauge theory description are given by eigenvalues of the adjoint scalars X^i and the K probes can be separated from the N targets by a VEV:

$$X^i = \begin{pmatrix} 0_{N \times N} & 0 \\ 0 & x I_{K \times K} \end{pmatrix} \quad (4)$$

We can relate x to the physical separation $\tilde{b}$ as $x = \tilde{b} \tilde{M}_S^2 = b R M_P^3$. The M-theory probe moves at a transverse velocity of v and the boost that produces the related system of Dp-branes rescales the velocity to $\tilde{v} = vR_S/R$. This velocity appears as the only nonvanishing component of the electric field in the gauge theory:

$$F_{0j} = \partial_0 X_j = \begin{pmatrix} 0_{N \times N} & 0 \\ 0 & f I_{K \times K} \end{pmatrix} \quad (5)$$

Here f is related to the physical velocity of the probe as $f = \tilde{v} \tilde{M}_S^2 = v R M_P^3$. The VEV in Eq. 4 spontaneously breaks the gauge group $U(N+K) \rightarrow U(N) \times U(K)$ and exchange of the resulting W-bosons produces the interaction. This can be studied efficiently by constructing the effective action as a function of b and v that results upon integrating out the W-boson.

The general form of the perturbative effective action resulting from this procedure can be understood using the dimensional analyses of [15, 8]:

$$L_{\text{eff}} = \frac{f^2}{2 g_{YM}^2} \left[1 + \sum_{L,m=1}^{\infty} c_{Lm} \left(\frac{g_{YM}^2 f^2}{x^{7-p}} \right)^L \left(\frac{f^2}{x^4} \right)^{m-L} \right] \quad (6)$$

Here L counts the number of Yang-Mills loops and $m+1$ counts the insertions of f^2 . In M theory variables $x = b R M_P^3$, $f = v R M_P^3$ and g_{YM}^2 is given in Eq. 3. At L loops, planar diagrams will have a total power of N and K adding to $L+1$. Nonplanar diagrams lower the total powers of N and K by an even number.² So at L loops the general N and K dependence is $N^{L-2n-q} K^{1+q}$ where $2n$ counts the degree of nonplanarity, and $q+1$ counts how many boundaries lie on the probe in a 't Hooft double line representation of the gauge theory diagram. The general form of the effective probe Lagrangian becomes:

$$L_{\text{eff}} = \frac{K v^2}{2 R} \left[1 + \sum_{L,m=1}^{\infty} \sum_{n=0}^{\lfloor (L-1)/2 \rfloor} \sum_{q=0}^{L-1-2n} d_{Lm}(n, q) K^q N^{2n-q} \right. \\ \left. \times \left(\frac{N v^2}{R^2 M_P^9 V_T b^{7-p}} \right)^L \left(\frac{v^2}{R^2 M_P^6 b^4} \right)^{m-L} \right] \quad (7)$$

(We have integrated this effective Lagrangian over the p spatial dimensions of the probe cancelling a factor of the dual torus volume.) The tree level (v^2) term is

²The total power of N and K is reduced by an even number because $U(N)$ diagrams in 't Hooft double line notation are orientable.

proportional to K because there are K probes and since the VEVs we have chosen are symmetric between the probes. Similarly, the factors of N arise because the probes interact with each brane in the target symmetrically. In principle, each factor of K and N should be replaced by traces over $U(K)$ and $U(N)$ gauge indices respectively and the effective action should be written covariantly in terms of F .

Eq. 7 is an analysis of the Feynman loop expansion of the gauge theory. However, there can also be contributions to the coefficients that are not captured by Feynman diagrams. Nonperturbative effects can also produce new terms that are simply not seen in perturbation theory. Indeed, we certainly know that they are necessary in considering graviton scattering in the presence of transverse p-tori for $p \geq 2$ [21, 22]. Also, we have not taken account of the elusive bound-state wavefunction governing the interacting clumps of branes in the gauge theory. Such bound state effects can easily add corrections to Eq. 7 with modified dependences on N and K . Analysis of such effects is very difficult, and so we are trying to learn as much as possible from general principles.

2.3 Scattering in supergravity

The limit where the scattering is described by gauge theory can in some cases also be treated accurately in supergravity. To determine when this is the case consider 11 dimensional supergravity compactified on a *spatial* circle of length R_S and a transverse T^p . This theory has plane wave solutions carrying N units of momentum in the compact direction:

$$ds^2 = dx^+ dx^- + dx_1^2 + \cdots + dx_9^2 + \tilde{D}(dx^+)^2 \quad (8)$$

$$\tilde{D} = \frac{c_p N}{(rM_P)^{7-p}(R_S M_P)^2 (V_T M_P^p)} \quad (9)$$

$$c_p = 4\pi^{\frac{p-1}{2}} \Gamma\left(\frac{7-p}{2}\right) \quad (10)$$

where $x^\pm = x^{11} \pm t$. Supergravity becomes inaccurate below the string scale so it may appear that this solution is only valid for $r \gg (R_S M_P^3)^{-1/2}$. This would be a very severe requirement because we would like to take $R_S \rightarrow 0$. Happily this conclusion is too hasty: Eq. 8 represents a solution on a spacelike circle $x_{11} \equiv x_{11} + R_S$. This metric can be rewritten as:

$$ds^2 = (-dt^2 + dx_{11}^2 + dx_1^2 + \cdots + dx_9^2) + \tilde{D}(dx_{11} + dt)^2 \quad (11)$$

$$\tilde{D} = \frac{c_p N}{(rM_P)^{7-p}(R_S M_P)^2 (V_T M_P^p)} \quad (12)$$

When the 11th circle is of length R_S we only expect supergravity to be valid at distances greater than the string length: $r \gg (R_S M_P^3)^{-1/2}$. However, Eq. 11 shows that the physical length squared of the circle is:

$$l^2 = \left(1 + \frac{c_p N}{(rM_P)^{7-p}(R_S M_P)^2 (V_T M_P^p)}\right) R_S^2 = R_S^2 + \frac{c_p N}{(rM_P)^{7-p} M_P^2 (V_T M_P^p)} \quad (13)$$

So, *independently* of R_S , in the large N limit (which we will take in the next section) the circle is much bigger than the 11 dimensional Planck length for any $r < \infty$. What is more, it can be checked that the curvature invariants scale inversely with N and so remain small in the limit. This strongly suggests that, for large N , the solution remains valid even in the $R_S \rightarrow 0$ limit. Essentially, the momentum along the circle exerts a pressure that decompactifies the solution allowing a valid treatment in 11 dimensional supergravity. From the 10 dimensional perspective, the solutions discussed above are D-particles or extremal black holes. We find that in the large N limit the throat region of the black hole extends far from the source and can be treated reliably in supergravity. Essentially, the local dilaton becomes large and so the solution should be considered 11 dimensional.

Now we use 11 dimensional Lorentz invariance to boost the compact direction by a rapidity β with $\sinh \beta = -R/\sqrt{2}R_S$. This converts the spacelike circle of length $x_{11} \equiv x_{11} + R_S$ into a nearly lightlike circle and the boosted metric (to leading order in R_S/R) is:

$$ds^2 = dx^+ dx^- + dx_1^2 + \cdots + dx_9^2 + D(dx^+)^2 \quad (14)$$

$$D = \frac{c_p N}{(rM_P)^{7-p}(RM_P)^2(V_T M_P^p)} \quad (15)$$

Since R_S has completely dropped out of the leading order boosted metric, *we can take R_S to zero freely without affecting the validity of the solution*. In other words, the boosted solution is valid for all $r \gg 1/M_P$. This is surprising because, as emphasized in [6, 19], the nearly lightlike compactified theory is related by a boost to the theory on a small spacelike circle where the string length would seem to be the relevant scale. In fact, as emphasized above, the supergravity solution on a spacelike circle has a much larger regime of validity than naively expected. A physical reason for the validity of the boosted solution at short distances is that x^+ , the putative compact “null” direction, is in fact *spacelike* for $r < \infty$. What is more, in the large N limit (which we will take in the next section), this *spacelike* circle is much larger than $1/M_P$ for any finite r . This strongly suggests that, for large N , we should be able to do reliable 11 dimensional supergravity calculations in this background.

In the limit of small R_S , the effective Lagrangian for a probe graviton carrying longitudinal momentum $P^+ = K/R$ has been constructed in [8]:

$$L_g = -\frac{K}{R} D^{-1} (\sqrt{1 - Dv^2} - 1) \quad (16)$$

3 Low energy, Large N limits

In this section we are going to take $N \rightarrow \infty$ limits while $R \rightarrow \infty$ in such a way that all energies and momenta are small. It is important that we take N and R to ∞ simultaneously and in a correlated fashion since we can always rescale R itself by a boost. We will consider two kinds of limits. First, following [6], we take $R_S \rightarrow 0$ before taking R and N to infinity. From the discussion of supergravity solutions in the previous section we find that this order of limits has a low energy supergravity

description. This result allows us to sum the leading N terms in the gauge theory effective action to all loops. Finally we consider a second limit in which R_S is held finite while N and R are taken to infinity. This is another decompactification limit of M theory. Assuming that the decompactification limit is unique leads to a specific conjecture for a relation between long and short distance string theory.

3.1 A supergravity limit (L1)

We begin by taking the $R_S \rightarrow 0$ limit of the nearly lightlike circle. Then we perform the following substitution on the parameters of the M theory:

$$L1 : \quad R \rightarrow \lambda R_0 \quad ; \quad N \rightarrow \lambda^2 N_0 \quad ; \quad K \rightarrow \lambda^2 K_0 \quad (17)$$

and take the scaling limit $\lambda \rightarrow \infty$. We will begin by arguing that this limit should be nonsingular and well described by the low-energy effective action of M theory.

Since N and K are getting very large there is some danger that $L1$ is a singular limit since the energies of the probe and target are very large. There are two energies relevant to our scattering processes: the energy stored in the longitudinal momentum which acts like a “mass”, and the energy in the transverse motion that acts like a “kinetic” energy. We start with the “kinetic” energy in the transverse motion of the probe. We can study this energy in a boost invariant way by examining the transverse momentum:

$$P_\perp^2 = P^+ P^- \sim \frac{K^2 v^2}{R^2} \quad (18)$$

In the decompactification limit, we expect the interactions between probe and target to be under control when the momentum density is small. The physical length of the compact direction scales proportionally to R , and so an appropriate requirement is:

$$\frac{P_\perp}{R} \sim \frac{Kv}{R^2} \ll M_P^2 \quad (19)$$

This relation is *independent* of the scaling limit $L1$ and simply requires a suitably small velocity.

Next we may worry that the mere fact that the longitudinal momenta of both probe and target ($P_p^+ = K/R$ and $P_t^+ = N/R$) are diverging will imply that a low energy description will break down due to strong gravitational effects. But we have already argued that the supergravity solution in Eq. 14 remains valid at small R_S . Evidence against the breakdown of this solution is provided by the smooth limit of the harmonic function D under the $L1$ scaling. So there is no reason to expect strong gravitational effects from the increasing “mass”, essentially because the “mass” density is not divergent in the decompactified $L1$ limit. Furthermore, as $N \rightarrow \infty$, we have argued that the compact direction becomes spacelike and large while curvature invariants remain small. This too suggests that gravitational effects are small.

Quantum corrections are immediately recognized as benign from the 11 dimensional point of view because all length scales are large. From the 10 dimensional point of view the local value of the dilaton is large close to the branes, suggesting a strong

coupling region in the interaction of D-branes and hence the gauge theory. However, as explained in [2], this apparent divergence is taken care of correctly by the gauge theory so that it is in fact the asymptotic dilaton that measures the magnitude of quantum corrections. The reason is that, at short distance, handles on diagrams with several boundaries are suppressed. In this way, it is precisely the circle of vanishing size at infinity, which is problematic for the 11 dimensional supergravity, that keeps the 10 dimensional theory under control.

Finally, consider the effective action for a supergravity probe given in Eq. 16. Expanding in powers of v gives a leading term $Kv^2/2R$. (Note that this is the same as the tree level term in the gauge theory effective action in Eq. 7.) In the $L1$ scaling limit, this “kinetic energy” term grows linearly in λ . This is a simple reflection of the fact that the total probe longitudinal momentum, which acts as an effective “mass”, grows linearly in the $L1$ limit. The growth with λ is just a consequence of the extensivity of the probe effective action and can be divided out. The condition that permits expansion of the supergravity effective action (Eq. 16) in powers of v is:

$$Dv^2 = \frac{c_p N v^2}{(rM_p)^{7-p} (RM_P)^2 (V_T M_P^p)} \ll 1 \quad (20)$$

Since D has a well defined limit under the $L1$ scaling, this requires small velocities and/or large distances in the scattering process.

Since we began by taking $R_S \rightarrow 0$, the discussion in Sec. 2 applies and M theory can be studied in the gauge theory description of low energy Dp-branes. So, to analyze the effect of the $L1$ limit on graviton scattering in M theory, we should study the effect it has on the gauge theory dynamics. Applying the scaling to the effective action in Eq. 7 we find that the tree level term scales linearly as λ . As discussed above, this overall scaling arises simply because of the extensivity of the probe action. After dividing out this overall scaling with λ , the individual terms in the rest of the Lagrangian scale as $\lambda^{2(L-2n-m)}$. So as $\lambda \rightarrow \infty$, the terms with $2n+m > L$ vanish, while the terms with $2n+m < L$ diverge. However, as discussed above, there are good reasons to believe that the $L1$ limit is a nonsingular limit of M theory. If so, we must conclude that the coefficients $d_{Lm}(n, q)$ vanish for $2n+m < L$. For example, for $n=0$ we are dealing with planar diagrams, and we are finding that all L loop planar diagrams with less than $2L+2$ velocity insertions must vanish. In other words, by assuming 11 dimensional Lorentz invariance and the plausible existence of the limit $L1$, we have shown an infinite number of non-renormalization theorems for the gauge theory effective action. For $p=0$, one of these theorems says that the one-loop v^4 term is perturbatively exact in the large N loop expansion. In fact, in this case, the vanishing of the coefficient of the two loop v^4 term ($L=2, m=1$) was checked in [23]. As we mentioned at the outset, our claims strictly concern terms of a form that can appear in the loop expansion. It is known, for example, that for $p=2$ there are non-perturbative contributions to the v^4 term [24] but these do not take the form of the terms in the loop expansion of Eq. 7.

The finite terms in the $L1$ limit are those that have $L=2n+m$. We insert this value of L and then relabel the parameter m as L . After an overall rescaling by $1/\lambda$

we find:

$$L_{L1} = \frac{K v^2}{2 R} \left[1 + \sum_{L=1}^{\infty} \sum_{q=0}^{L-1} d_{LL}(0, q) K^q N^{-q} \left(\frac{N v^2}{R^2 M_p^9 V_T b^{7-p}} \right)^L (1 + S_{L1}) \right] \quad (21)$$

Here S_{L1} is a series of corrections suppressed by inverse powers of M_P^6 :

$$S_{L1} = \sum_{n=1}^{\infty} f_{Lq}(n) \left(\frac{v^2}{R^2 M_p^9 V_T b^{7-p}} \right)^{2n} \left(\frac{v^2}{R^2 M_p^6 b^4} \right)^{-2n} = \sum_{n=1}^{\infty} f_{Lq}(n) \left(\frac{1}{M_P^3 V_T b^{3-p}} \right)^{2n} \quad (22)$$

where $f_{Lq}(n)$ are constant coefficients. We want to compare this limiting action to low energy, decompactified M-theory. Since M_P is finite in our limit, we expect corrections to scattering amplitudes that arise from subleading corrections to the M theory low energy effective action. Russo and Tseytlin have shown that the curvature corrections to M-theory that derive from perturbative corrections in string theory are of the form $\mathcal{R}^{3k+1}/M_P^{6k-9}$ [25] where $\mathcal{R}$ is the Riemann tensor. In our limit of the gauge theory, the subleading terms in M_P are in the series S_{L1} which collects terms arising from nonplanar diagrams. It is pleasing that S_{L1} is a series in $1/M_P^6$ in accord with our expectation from Russo and Tseytlin. The terms in the effective action that have $n = 0$ (and come from planar diagrams) should be identified with the leading term in the M theory low energy effective action - 11 dimensional supergravity. It is interesting that the quantum corrections in M-theory should appear in “diagonal” bands ($L = 2n + m$) whereas the quantum expansion in the gauge theory is “vertical” (in L).

To study the planar diagrams with $n = 0$ note that the effective loop expansion parameter in Eq. 21 is:

$$g_{\text{eff}}^2 = \frac{N v^2}{(b M_p)^{7-p} (R M_P)^2 (V_T M_P^p)} \quad (23)$$

The condition that this should be small, so that the perturbative treatment is valid, is precisely the same as the condition in Eq. 20 that permitted expansion of the supergravity effective action in Eq. 16 in an eikonal type series in powers of velocity. For $n = 0$, the finite terms in the $L1$ limit are, up to coefficients, an expansion in $D v^2$ where D is the harmonic function in the supergravity solution of Eq. 14. This means that the $n = 0$ terms in the gauge theory action that scale as $N^L K$ coincide (up to the undetermined coefficients $d_{LL}(0, 0)$) with the expansion in powers of velocity of the supergravity effective action in Eq. 16. In the next section we will take a different large N limit that isolates just these terms. In the $L1$ limit, the total longitudinal momentum of the probe diverges and so back-reaction effects are important. The terms with higher powers of K can be interpreted as arising from back-reaction of the probe gravitons. The supergravity probe action in Eq. 16 was derived in the absence of back-reaction - hence we cannot hope to compare it directly to the gauge theory in the $L1$ limit.

3.2 A supergravity probe limit (L2)

In the previous section we found that the terms with $L = m$ and $n = 0$ in the gauge theory effective action were associated with supergravity. In this section we will take another limit where back-reaction of the probe is suppressed so that the supergravity effective action in Eq. 16 can be directly compared to the gauge theory action. The new limit is defined by the substitutions:

$$L2 : \quad R \rightarrow \lambda R_0 \quad ; \quad N \rightarrow \lambda^2 N_0 \quad ; \quad K \rightarrow \lambda K_0 \quad (24)$$

followed by $\lambda \rightarrow \infty$. Again, $R_S \rightarrow 0$ before these limits are taken. Following the analysis of the previous section, the longitudinal momentum density of the target $P_t^+/R \sim N/R^2$ is fixed in the $L2$ decompactification limit. However, the probe longitudinal momentum density $P_p^+/R \sim K/R^2$ and the transverse momentum density $P_\perp/R \sim K v^2$ vanish as $\lambda \rightarrow \infty$ although the total probe momenta $P_p^+ \sim K/R$ and $P_\perp \sim K v^2/R$ remain finite. We have already argued that the $L1$ limit should be a nonsingular limit of M theory. The same discussion applies here, with the added statement that the vanishing energy density of the probe implies that it can be treated as moving in the background produced by the target without any back-reaction.

Examining the effect of the $L2$ scaling limit on the gauge theory effective action in Eq. 7 shows that the tree level term $K v^2/R$ remains finite. This is because, unlike the $L1$ limit, the total energy of the probe remains finite. The general term in the effective action scales as $\lambda^{2(L-2n-m)-q}$. Again, as with the $L1$ limit, we have good reasons to assume that the $L2$ limit is nonsingular and so all divergent terms must in fact have vanishing coefficients. The finite terms satisfy $2n + m = L - q/2$. However, the $L1$ limit has already shown that terms with $2n + m < L$ must vanish. So only the terms with $q = 0$ survive the $L2$ limit. We are left with the following finite terms in the effective action:

$$L_{L2} = \frac{K v^2}{2R} \left[1 + \sum_{L=1}^{\infty} d_{LL}(0,0) \left(\frac{N v^2}{R^2 M_P^9 V_T b^{7-p}} \right)^L (1 + S_{L2}) \right] \quad (25)$$

Here S_{L2} is a series of corrections suppressed by inverse powers of M_P^6 :

$$S_{L2} = \sum_{n=1}^{\infty} f_{L0}(n) \left(\frac{1}{M_P^3 V_T b^{3-p}} \right)^{2n} \quad (26)$$

The series S_{L2} collects the contributions from nonplanar diagrams that survive the $L2$ limit. The terms in S_{L2} are suppressed by powers of $1/M_P^6$ which matches our expectations for the corrections to the low-energy effective action of M theory in the decompactified limit [25]. The $n = 0$ terms arise from planar diagrams and should be compared to supergravity.

Therefore we compare the planar ($n = 0$) diagrams that survive the $L2$ limit to the supergravity probe effective action in Eq. 16. Evidently, the gauge theory loop expansion coincides up to the unknown coefficients $d_{LL}(0,0)$ with the velocity expansion of the supergravity effective Lagrangian. In the limit of parameters we are

studying, as we have argued above, both the gauge theory and the 11 dimensional supergravity analyses appear to be valid. So the probe Lagrangian in Eq. 16 is expected to be the summation of planar diagrams that survive the $L2$ limit. In other words, we conclude that the summation to all loops of the leading N planar diagrams in the gauge theory effective action gives a Born-Infeld action of the form:

$$L = -\frac{K}{R}D^{-1}(\sqrt{1-Dv^2}-1) \quad (27)$$

One way of stating this result is that the leading large N terms in the gauge theory effective action are controlled by a symmetry - 11 dimensional Lorentz invariance - that causes them to resum to a Born-Infeld form.

Independent Evidence for Born-Infeld resummation: The same $U(N+K)$ gauge theories studied above can be derived by keeping R_S fixed and examining the short distance dynamics of Dp-branes where the probes have energies $E \sim (1/2)Kv^2R_S/R^2$ with $R \gg R_S$. It is well known that for any p this low-energy, short-distance dynamics is governed by $p+1$ dimensional gauge theory [1, 2]. The gauge theory should be understood here as merely a low-energy effective description of the branes - in previous sections, the $R_S \rightarrow 0$ limit made gauge theory the exact theory for $p \leq 3$.

In this alternative scenario, the analysis of the gauge theory effective action for scattering of K probe from N target branes proceeds exactly as in Sec. 2. As before, we separate the probes spatially from the target by a distance $\tilde{b}$, and we give the probes a velocity represented by the electric field Eq. 5. In addition, we turn on constant magnetic fluxes restricted to live in $U(1)^K \in U(K)$ in order to avoid difficulties when the fluxes do not commute. Then, for any p , ignoring cutoff dependent terms, the effective action governing scattering of K probes from the target will be given by Eq. 7 with the following modifications. Each v^{2m+2} is replaced by a Lorentz scalar constructed from $(2m+2)$ factors of F , the flux on the probe. Also, each factor of K is replaced by a trace over the probe gauge fields. When there is more than one trace it is hard to know how to distribute them over the factors of F and the relative coefficients between the different Lorentz scalars at $O(F^{2m+2})$ should be determined. (These issues have been discussed in [9, 10].)

The effective actions for various p all arise as low-energy limits of open-string theory. Since T-duality is a symmetry of open-string theory, we expect that effective actions for different p should be related to each other by dimensional reduction [9]. We will use this expectation along with supersymmetry and existing calculations to derive the leading N gauge theory effective action for any p up to 2-loops. The tree level Lagrangian (after integrating the constant fluxes over the Dp-brane worldvolume) is:

$$L_0 = \frac{\text{Tr}(F^2)}{4R} \quad (28)$$

We are interested in the leading N planar diagrams which, as discussed in previous sections, have $n = 0$ and are proportional to N^L at L loops. After integrating over

the p spatial dimensions, the general form of the terms with $(2L + 2)$ factors of F at 1 and 2 loops is:

$$L_1 = (1/R)\text{Tr}(\alpha F^4 + \beta (F^2)^2) \quad (29)$$

$$L_2 = (1/R)\text{Tr}(\gamma F^6 + \sigma F^4 F^2 + \tau (F^2)^3) \quad (30)$$

(We use the convention that F^{2n} has a cyclic trace on Lorentz indices.)

When F is a pure electric field (transverse velocity) as in Eq. 5, the value of L_1 and L_2 have been computed in [2, 8]. When $p = 4$ and F is self-dual, supersymmetry is unbroken and there is no force between probe and target. So in this case L_1 and L_2 vanish. Finally, if $p = 4$, and we turn on a self-dual magnetic flux as well as the transverse velocity in Eq. 5, we expect a metric on moduli space that is exact at one loop [2]. This gives another vanishing condition on L_2 .

Altogether we have five equations for five unknowns which can be solved to give:³

$$L_1 = \frac{D}{R} \text{Tr} \left(\frac{F^4}{8} - \frac{(F^2)^2}{32} \right) \quad (31)$$

$$L_2 = \frac{D^2}{R} \text{Tr} \left(\frac{F^6}{12} - \frac{F^4 F^2}{32} + \frac{(F^2)^3}{384} \right) \quad (32)$$

Here D is the harmonic function in Eq. 15. With some work it can be shown that $L_0 + L_1 + L_2$ is the expansion to $O(F^6)$ of the Born-Infeld like Lagrangian:

$$L_{BI} = -(1/R) D^{-1} \text{Tr} \left(\sqrt{\det(\eta_{\mu\nu} I - D^{1/2} F_{\mu\nu})} - I \right) \quad (33)$$

The determinant is over the Lorentz indices and I is the $U(K)$ identity. It is rather non-trivial that the one and two loop effective Lagrangians fit the expansion of the Born-Infeld formula. Previously, when F simply represented a transverse velocity, we derived the same Born-Infeld form for the leading N planar diagrams that survive the $L2$ limit. (In that case, $\det(\eta_{\mu\nu} I - D^{1/2} F_{\mu\nu}) = (1 - Dv^2)I$.) That derivation used 11 dimensional Lorentz invariance and used the plausible existence of certain decompactification limits of M theory. Here we have made the rather different assumption that gauge theory effective actions are related by T-duality. As a consequence of this assumption, supersymmetry and existing calculations provide significant evidence for the Born-Infeld resummation of the leading N planar diagrams studied in previous $L2$ limit. This can be seen as partial evidence for the existence of the decompactification limit assumed in previous sections.

3.3 Another large N decompactification limit (L3)

In this section we will analyze a large N limit of a different sort from before. Let us consider nearly null compactified M-theory specified by N, R and $\epsilon = \frac{R_s}{R}$, a small parameter measuring the almost “light-like-ness”. We will assume that if a decompactified limit

$$N, R \rightarrow \infty, \epsilon \rightarrow 0 \quad (34)$$

³The identity $F^{2n} = (1/4^{2n-1})(F^2)^n$ for self-dual fields is very useful in showing this.




Figure 1: Two possible decompactification limits

exists and is well defined, it is unique. This is a rather strong assumption, but it seems reasonable in that if there are a number of consistent limits then there would seem to be many noncompact 11-dimensional vacua. So let us make this assumption and see what conclusions we are led to. We will then see some evidence for it.

In particular, let us examine two separate ways of taking the limit in Eq. 34. The first limit is the one that we have been studying thus far. We start in the lower left corner of the triangle in Fig. 1, at some given values of N , R and ϵ . We then follow the horizontal line in figure by taking the $\epsilon \rightarrow 0$ limit following Seiberg [6]. In this limit N and R are finite. As shown by Seiberg, this limit of M-theory, at the lower right corner of the figure, is related by a boost to weakly coupled strings in the gauge theory limit. The vertical line then takes $N, R \rightarrow \infty$ following the earlier L1 or L2 limits which, we argued, led to decompactified M theory. Thus, finally, the $N = \infty$ gauge theory describes decompactified 11-dimensional physics.

The second limit that we can take is the one sketched along the diagonal of the figure. Here we simultaneously take $N, R \rightarrow \infty$ and $\epsilon \rightarrow 0$ by keeping R_s fixed. We will refer to this as the “diagonal limit”. As before, the precise way in which we take $N \rightarrow \infty$ depends on what we want to study. In particular one could consider either of the L1 or L2 limits. After repeating Seiberg’s boost and rescaling of units as explained in Sec. 2, we find ourselves in string theory. This time we are in a regime where $\tilde{g}_S = (R_S M_P)^{3/2}$ is fixed and $\tilde{M}_S^2 = \frac{R^2 M_P^3}{R_S} = \frac{R_S M_P^3}{\epsilon^2}$. The velocities are $\tilde{v} = v\epsilon$ and the transverse separations $\tilde{b} = b\epsilon$. Unlike the previous $R_S \rightarrow 0$ limit, the physical separations in string units $\tilde{b}\tilde{M}_S = b(R_S M_P^3)^{1/2} = bM_P \tilde{g}_S^{1/3}$ are fixed. So we are not in the gauge theory limit of string theory - the limit along the diagonal in Fig. 1 is related to a rather different limit of string theory. If this also describes decompactified M-theory, then an equality of limits would imply that these two limits of *string theory* are the same. At first sight this seems completely unlikely and therefore to undercut our initial assumption. But let us examine it a little more.

We work in the domain where the decompactified theory would be adequately described by 11-dimensional supergravity which we can then compare with our second limit of string theory. To this end we take $bM_P \gg 1$ so that for fixed (small) $\tilde{g}_S$ we have $\tilde{b}\tilde{M}_S \gg 1$. This takes us to a domain the string theory is in a 10-dimensional supergravity limit. For concreteness, consider the process of two graviton scattering in the probe limit (L2) Eq. 24. In string theory we are then computing the effective action for the probe D-particle in the presence of the background fields produced by the target D-particle. For small string coupling this is given by the disc with Dirichlet boundary conditions in the presence of background fields.

$$\begin{aligned} S_{\text{eff}} &= \frac{K\tilde{M}_S}{\tilde{g}_S} \int dt \{e^{-\phi}(\det G)^{1/2} - A\} \\ &= \frac{K\tilde{M}_S}{\tilde{g}_S} \int dt f^{-1}(\sqrt{1 - f\tilde{v}^2} - 1) \end{aligned} \quad (35)$$

where

$$f = 1 + \frac{c\tilde{g}_S N}{(\tilde{r}\tilde{M}_S)^7} \quad (36)$$

As we take the limit $\epsilon \rightarrow 0$ (keeping R_s fixed in the “diagonal limit” in the figure), we see that the disc effective action has a well defined limit:

$$S_{\text{eff}} = - \int dt \frac{K}{R} D^{-1}(\sqrt{1 - Dv^2} - 1) \quad (37)$$

$$D = \frac{c_0 N}{(rM_P)^7 (RM_P)^2} \quad (38)$$

Note that the leading 1 in Eq. 36 has dropped away in the limit. We are left with precisely the same action as the *11-dimensional* lightcone supergravity effective action that we had in Eq. 16. This is what we had earlier claimed would be the leading term to survive in the gauge theory limit of string theory in the L2 limit. This is then some evidence that our conjecture of equality of different limits of string theory is not altogether meaningless. Note that all dependence on R_s , which was fixed, has gone away. This is essentially like being in the near horizon regime of black holes in string theory. The simultaneous validity of large N gauge theory and supergravity in this domain has already been proposed [13]. Our argument about the existence and equality of various decompactification limits is a sort of a generalisation of this to general string theory processes.

We can make further testable statements. In the second string limit, there are corrections beyond the disc diagram in a background field. These are of two kinds. There are the diagrams with one boundary and any number of handles attached. Therefore they come with only one power of K and hence are corrections to supergravity but nevertheless in the probe limit. There are corresponding candidate M_P^6 corrections in the gauge theory from below the diagonal. It will be interesting to verify that, say, the disc with one handle attached, in the presence of the target fields, gives the same contribution as non-planar gauge theory diagrams from one below the diagonal. If we go beyond the probe approximation one can consider string diagrams with extra

boundaries. Since these have higher powers of K , they take into account the effect of the probe on the geometry. Indeed, our conjecture leads us to expect that the cylinder diagram with both boundaries on the probe, in the background produced by the source should reproduce the gauge theory planar diagrams that are of order K^2 and survive the L1 limit.

4 Discussion

Let us review the results we have obtained. We considered M theory on a nearly lightlike circle and a transverse torus with N units of longitudinal momentum. We defined three kinds of large N limits of this system, and argued that they were nonsingular. Furthermore, we argued that 11 dimensional supergravity is valid in these limits because the effect of the momentum in the supergravity solution is to make the circle spacelike and big. The M theoretic description of graviton scattering then allowed us to infer properties of gauge theory such as an infinite set of non-renormalisation theorems. The compatibility and existence of the third of our limits led to a conjectured relation within string theory between two very different regimes. This had surprising, but testable, consequences.

The large N limits that we have taken are not conventional 't Hooft limits. In fact, in ordinary gauge theories, if we take the large N limit keeping the gauge coupling finite, the Feynman diagram expansion is wildly divergent. Why then do we expect our limits to be well defined? Our argument has been indirect, relying on the well-behaved nature of supergravity. We can turn this around and say that if supersymmetric Yang-Mills has vanishing coefficients for the infinite class of terms that we identified, then the unconventional large N limits we defined are finite.

The argument that supergravity is valid at short distances in the large N limit is also an important ingredient in other recent works [26, 27]. In the latter reference it is argued that superconformal invariance causes the leading N terms in the effective action of $3 + 1$ dimensional Yang-Mills to be of the Born-Infeld form. It would be interesting to understand the relation between our arguments and those of [27].

Several recent works [14, 15, 16, 17, 18, 20] have reported disagreements between finite N calculations in gauge theory and supergravity. There is no contradiction between these results and ours because we are dealing with the large N limit. We will, however, comment briefly on the reported finite N discrepancy in the simplest generalisation of our setting - three graviton scattering [18]. Consider two gravitons at a separation of r and another, much further away at a distance R . According to perturbative supergravity in 11 dimensions, the interaction includes a term of the form:

$$L_{3g} \sim \kappa^4 N_1 N_2 N_3 \frac{(v_1 - v_2)^2 (v_2 - v_3)^2 (v_1 - v_3)^2}{r^7 R^7} \quad (39)$$

It was argued in [18] that such a term cannot arise in finite N gauge theory. We have interpreted this scattering problem in 10 dimensions as the interaction of 3 D-particles that are almost at rest. In the gauge theory the relevant problem involves the symmetry breaking $SU(3) \rightarrow U(1)^2$. This can be analyzed in two parts, by first

considering $SU(3) \rightarrow U(2)$, which leads to an effective action with vertices of the v^4 form. This modified effective $U(2)$ Lagrangian has mildly broken supersymmetry, so that the subsequent step $U(2) \rightarrow U(1)^2$ gives rise to a v^2 force instead of v^4 . In this way we do arrive at terms of the general form $\kappa^4 \frac{N_1 N_2 N_3 v^6}{r^7 R^7}$. The remaining issue is whether we reproduce the precise kinematic factor in Eq. 39.

The crucial second step that gives a v^2 force can be interpreted in supergravity as the interaction between a D-particle and a near-extremal D-particle [29]. We have exploited this to establish a quantitative agreement between supergravity and gauge theory – a specific contribution to the 3 D-particle interaction. This contribution from gauge theory already serves to indicate the source of the tension between supergravity and M(atrix) theory in [18]: the gauge theory does not exhibit manifest Lorentz invariance in 11 dimensions. There *are* contributions of roughly the right form, but they do not naturally organize themselves into Lorentz invariants, and so all terms must be calculated before we can compare with Eq. 39.⁴

The authors of the various papers pointing out difficulties in matching gravity from the M(atrix) model have suggested that the problems may disappear in the large N limit. In this paper we have defined several large N limits that appear to have sensible low energy supergravity descriptions in M theory. As such, this is the arena in which the M(atrix) model should confront supergravity.

Acknowledgments: We would like to thank Shanta de Alwis, Shyamoli Chaudhuri, Mike Dine, Eric Gimon, David Kastor, Albion Lawrence, David Lowe, Juan Maldacena, Samir Mathur, Djordje Minic, Hirosi Ooguri, Arvind Rajaraman, Sanjaye Ramgoolam, Nati Seiberg, Andy Strominger and Jennie Traschen for useful discussions. V.B. thanks the University of Massachusetts at Amherst for hospitality and facilities. F.L. thanks MIT for hospitality while this work was completed. V.B. is supported by the Harvard Society of Fellows and by the NSF grant NSF-Phy-91-18167. R.G. is supported in part by DOE grant DOE-91-ER40618. F.L. is supported in part by DOE grant FG02-95ER40893.

References

- [1] E. Witten. *Nucl. Phys. B*, 460:335–350, 1996. hep-th/9510135.
- [2] M. Douglas, D. Kabat, P. Pouliot, and S. Shenker. *Nucl. Phys. B*, 485:85–127, 1997. hep-th/9608024.
- [3] T. Banks, W. Fischler, S. Shenker, and L. Susskind. *Phys. Rev. D*, 55:5112, 1997. hep-th/9610043.
- [4] L. Susskind. hep-th/9704080.
- [5] R. Dijkgraaf, E. Verlinde, and H. Verlinde. *Nucl. Phys. B*, 500:43–61, 1997. hep-th/9703030.

⁴Work in progress.

- [6] N. Seiberg. *Phys. Rev. Lett.*, 79:3577–3580, 1997. hep-th/9710009.
- [7] A. Sen. hep-th/9709220.
- [8] K. Becker, M. Becker, J. Polchinski, and A. Tseytlin. hep-th/9706072.
- [9] I. Chepelev and A.A. Tseytlin. hep-th/9709087.
- [10] E. Keski-Vakkuri and P. Kraus. hep-th/9709122.
- [11] P. Berglund and D. Minic. hep-th/9708063.
- [12] S. de Alwis. hep-th/9710219.
- [13] M. Douglas, J. Polchinski, and A. Strominger. hep-th/9703031.
- [14] M.R. Douglas, H. Ooguri, and S.H. Shenker. *Phys. Lett. B*, 402:36–42, 1997. hep-th/9702203.
- [15] O. Ganor, R. Gopakumar, and S. Ramgoolam. hep-th/9705188.
- [16] J.R. David, A. Dhar and G. Mandal. *Phys.Lett.B*415:135-143,1997.
- [17] M.R. Douglas and H. Ooguri. hep-th/9710178.
- [18] M. Dine and A. Rajaraman. hep-th/9710174.
- [19] S. Hellerman and J. Polchinski. hep-th/9711037.
- [20] E. Keski-Vakkuri and P. Kraus. hep-th/9712013.
- [21] J. Polchinski and P. Pouliot. *Phys. Rev. D*, 56:6601–6606, 1997. hep-th/9704029.
- [22] T. Banks, W. Fischler, N. Seiberg, and L. Susskind. *Phys. Lett. B*, 408:111–116, 1997. hep-th/9705190.
- [23] K. Becker and M. Becker. *Nucl. Phys. B*, 506:48–60, 1997. hep-th/9705091.
- [24] M. Dine and N. Seiberg. *Phys.Lett.*, B409:239–244, 1997. hep-th/9705057.
- [25] J. Russo and A. Tseytlin. hep-th/9707134.
- [26] J. Maldacena and A. Strominger. hep-th/9710014.
- [27] J. Maldacena. hep-th/9711200.
- [28] M. Serone. hep-th/9711031.
- [29] J. Maldacena. hep-th/9705053.